



New in vitro colonic fermentation model for Salmonella infection in the child gut

Gwenaëlle Le Blay, Julia Rytka, Annina Zihler, Christophe Lacroix

► To cite this version:

Gwenaëlle Le Blay, Julia Rytka, Annina Zihler, Christophe Lacroix. New in vitro colonic fermentation model for Salmonella infection in the child gut. FEMS Microbiology Ecology, 2009, 67 (2), pp.198-207. 10.1111/j.1574-6941.2008.00625.x . hal-00558484

HAL Id: hal-00558484

<https://hal.univ-brest.fr/hal-00558484>

Submitted on 21 Jan 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

New *in vitro* colonic fermentation model for *Salmonella* infection in children gut

Gwenaëlle Le Blay^{1,2}, Julia Rytka¹, Annina Zihler¹ and Christophe Lacroix¹

¹Laboratory of Food Biotechnology, Institute of Food Science and Nutrition, ETH Zurich, 8092 Zurich, Switzerland, ²Laboratoire Universitaire de Biodiversité et Ecologie Microbienne, ESMISAB, Université Européenne de Bretagne - Université de Brest, Technopôle Brest-Iroise, 29280 Plouzané, France.

Keywords: child / immobilized cells / intestinal microbiota / *in vitro* continuous fermentation model / *Salmonella* serovar Typhimurium

Running title: *In vitro* salmonellosis model

Corresponding author: Christophe Lacroix, ETH-Zurich, Institute of Food Science and Nutrition, Schmelzbergstrasse 7, LFV C20, 8092 Zürich, Switzerland. Tel.: +41 44 632 48 67; fax: +41 44 632 14 03; e-mail: christophe.lacroix@ilw.agrl.ethz.ch

Abstract

In this study, a new *in vitro* continuous colonic fermentation model of *Salmonella* infection with immobilized child fecal microbiota and *Salmonella* serovar Typhimurium was developed for the proximal colon. This model was then used to test the effects of two amoxicillin concentrations (90 and 180 mg day⁻¹) on microbial composition and metabolism of the gut microbiota and on *Salmonella* serovar Typhimurium during a 43-day fermentation. Addition of gel beads (2%, v/v) colonized with *Salmonella* serovar Typhimurium in the reactor resulted in a high and stable *Salmonella* concentration (log 7.5 cell number mL⁻¹) in effluent samples and a concomitant increase of *Enterobacteriaceae*, *Clostridium coccoides*-*Eubacterium rectale* and *Atopobium* populations and a decrease of bifidobacteria. During amoxicillin treatments, *Salmonella* concentrations decreased while microbial balance and activity were modified in agreement with *in vivo* data with a marked decrease of *Clostridium coccoides*-*Eubacterium rectale* and an increase in *Enterobacteriaceae*. After interruption of antibiotic addition, *Salmonella* concentration increased again to reach values comparable to that measured before antibiotic treatments, showing that our model can be used to simulate *Salmonella* shedding in children as observed *in vivo*. This *in vitro* model could be a useful tool for developing and testing new antimicrobials against enteropathogens.

45 Introduction

Salmonellosis is one of the most common and widely distributed foodborne diseases worldwide. It is associated with two types of symptoms, caused by different serovars of *Salmonella enterica* subsp. *enterica*: typhoid fever (mainly caused by serovar Typhi, Paratyphi and Sendai) which is more common in developing countries; and gastroenteritis (mainly caused by serovar Typhimurium and Enteritidis) also encountered in developed countries (Coburn *et al.*, 2007). In Europe, 176,395 cases of salmonellosis infections (i.e. 38 for 100 000 habitants) have been declared in 2005, with a major proportion (20%) of young children less than 5 years of age (The European Food Safety Authority & European Center for Disease Prevention and Control, 2006). In children, dehydration associated with diarrhea can become severe and life-threatening (Rosanova *et al.*, 2002), therefore effective antimicrobials are essential drugs for treatment. Antimicrobials most widely regarded as optimal for treating diarrheal diseases caused by *Salmonella* serovar Typhimurium in children are third-generation cephalosporins, because quinolones and fluoroquinolones are generally not recommended due to their toxicity on immature cartilage and the possible emergence of resistant pathogens (Schaad, 2005). The earlier drugs, chloramphenicol, ampicillin, amoxicillin and trimethoprim-sulfamethoxazole, are also used as alternatives (Frye & Fedorka-Cray, 2007; World Health Organization, 2008). However, the emergence of *Salmonella* isolates with multiple-drug resistance urges the need for alternatives to antibiotherapies (White *et al.*, 2001; World Health Organization, 2008). Valuable models of *Salmonella* infection are therefore needed to develop and test new treatments, in particular for young human populations. To our knowledge, there is currently no suitable model to test the effects of new antimicrobials on both enteropathogens and intestinal bacteria. Animal models, such as bovine or streptomycin-pretreated mice models for *Salmonella* infection do not reproduce the human intestinal microbiota and are dedicated for studying host-pathogen interactions

70 (Hapfelmeier & Hardt, 2005). Moreover, *in vivo* studies are difficult to perform due to cost, ethical problems and high interindividual variations. By contrast, *in vitro* models are much less expensive, simpler to handle and could be a good alternative for testing new antimicrobial treatments in a first screening phase; but current models are not suitable for intestinal infections. The continuous culture models for human intestinal microbiota are
75 mainly based on the original model of Macfarlane *et al.* (1998) and inoculated with diluted feces. These systems have several limitations due to the planktonic state of bacterial populations, leading to limited microbial stability and cell density compared to the colon; they are also not suited for long-term experiments with enteropathogens since exogenous bacteria are rapidly washed out of the system (Blake *et al.*, 2003; Carman *et al.*, 2004; Carman &
80 Woodburn, 2001; Payne *et al.*, 2003). Recently, we developed and validated a new model of infant and adult colonic fermentation with fecal microbiota immobilized in gel beads in anaerobic continuous-flow cultures (Cinquin *et al.*, 2006a; Cinquin *et al.*, 2006b; Cleusix *et al.*, 2008; Macfarlane & Macfarlane, 2007). This new model has conditions that are more akin to that of the intestinal system and has the following characteristics: bacteria growing in
85 biofilm structures; high cell density in gel beads and in reactor effluents (up to 10^{11} cells mL⁻¹ or g⁻¹); high stability and reactivity to changing conditions of the intestinal fermentation; good protection of sensitive bacteria from shear and oxygen stresses; and prevention of washout and loss of less competitive bacteria.

In this study we developed an *in vitro* model of intestinal fermentation with immobilized
90 feces simulating intestinal *Salmonella* infections and long-term shedding in children. We hypothesized that addition of colonized beads with *Salmonella* serovar Typhimurium in the intestinal fermentation model containing an immobilized child microbiota would cause a stable infection of *Salmonella* in gel beads and in effluent samples. The effect of two antibiotic treatments on *Salmonella* serovar Typhimurium, as well as on the main bacterial

95 populations and metabolism of child's microbiota were tested during the same continuous culture and compared with *in vivo* data to validate the model.

Materials and methods

Bacterial strain

Salmonella enterica subsp. *enterica* serovar Typhimurium M557 (*sseD::aphT ΔinvG*)

100 (*S. Typhimurium*) was supplied by Prof. W. Hardt (Institute of Microbiology, ETH, Zurich, Switzerland). Sensitive to amoxicillin, this strain is a low virulent derivative of *S. Typhimurium* wild-type strain SL1344 lacking SPI-1 effector proteins (Hapfelmeier *et al.*, 2004). It was routinely cultivated in tryptone soya broth (TSB, Oxoid, Basel, Switzerland) overnight at 37 °C in aerobiosis.

105 *Antibiotic*

In human practice, Clamoxyl®/-RC (GlaxoSmithKline) containing amoxicillin as active compound, is a moderate-spectrum β-lactam antibiotic that can be used to treat salmonellosis in children (25-75 mg kg⁻¹ day⁻¹) (Moulin *et al.*, 2003; Fachinformation des Arzneimittel-Kompodium der Schweiz®, 2007). It is bactericidal against a wide range of Gram-positive and Gram-negative bacteria, including sensitive *Salmonella* with minimum inhibitory concentrations (MIC) ranging from 1 to 5 µg mL⁻¹ for 90% of tested strains (Fachinformation des Arzneimittel-Kompodium der Schweiz®, 2007). For the experiment, pure amoxicillin (Sigma-Aldrich, Buchs, Switzerland) was used.

Feces collection and bacterial immobilization

115 The fecal sample used for immobilization was collected from a healthy 2 year-old child, who had not received antibiotics at least 3 months before the experiment. The fecal sample was maintained in anaerobiosis and immobilized in 1-2 mm gel beads composed of gellan (2.5%,

w/v) and xanthan (0.25%, w/v) gums and sodium citrate (0.2%, w/v), as already described

(Cleusix *et al.*, 2008). Gel beads (60 mL) were then transferred into a stirred glass reactor

120 (Sixfors, Infors, Bottmingen, Switzerland) with 140 mL of fresh nutritive medium simulating a child chyme (presented below). The entire process was completed under anaerobic conditions within 3 h after defecation.

S. Typhimurium immobilization was done one day before reactor inoculation using the same procedure as for fecal samples, but in aerobiosis under a sterile bench. *Salmonella* beads

125 (10 mL) were colonized overnight in 200 mL TSB at 37 °C in aerobiosis without pH-control.

A fresh bead sample (0.5 g) was used to inoculate the reactor on day 11 and the rest was stored frozen (0.5 g aliquots) in 20% glycerol at -80 °C.

Nutritive medium

The nutritive medium used to feed the reactor was similar to that previously described by

130 Macfarlane *et al.* (1998) for simulating an adult ileal chyme, with one modification; the bile salt concentration was reduced from 0.4 to 0.05 g L⁻¹ to reproduce the ileal chyme of a young child. A solution of vitamins described by Michel *et al.* (1998) and sterilized by filtration (Minisart 0.2 µm, Sartorius, Göttingen, Germany) was added (0.5 mL l⁻¹) separately to the autoclaved (15 min, 121 °C) medium.

135 ***Experimental setup and sampling***

A single-stage reactor based on the model described by Cinquin *et al.* (2004) was used to mimic the microbial ecosystem of a child's proximal colon. Batch fermentations were first carried out to colonize the fecal beads for 2 days. During colonization, the nutritive medium was aseptically replaced by fresh nutritive medium every 12 h. Temperature (37 °C) was

140 automatically controlled and pH was maintained at 5.7 by adding NaOH (5N). Anaerobic conditions were maintained during the whole fermentation by a continuous flow of pure CO₂ in the headspace. The continuous fermentation was carried out in the same reactor connected

to a stirred feedstock vessel containing sterile nutritive medium continuously flushed with CO₂ and maintained at 4 °C and to an effluent receiving vessel. Continuous medium feeding was carried out using peristaltic pumps (Reglo analog, Ismatec, Glattbrugg, Switzerland) delivering a feed flow rate of 40 mL h⁻¹ for a mean retention time of 5 h. This time was used to simulate the residence time in a child proximal colon, with a pH of 5.7 (Fallingborg *et al*, 1990).

The 43-day continuous fermentation was divided into 6 periods of 5 to 9 days (Figure 1). First the system was stabilized (STyphi 0; days 3-10), then 0.5 g of beads colonized with *S. Typhimurium* (days 11 and 13) were added to the system followed by a second stabilization period (STyphi I; days 14-20) and two antibiotic treatments (ATB I [214 µg mL⁻¹ thrice a day]; days 21-25 and ATB II [428 µg mL⁻¹ thrice a day]; days 35-39) intercalated with a third stabilization period without antibiotic (STyphi II, day 26-34). Amoxicillin was added directly in the reactor three times per day (at 9 a.m., 2 p.m. and 6 p.m.) to reach total concentrations of 90 mg day⁻¹ (ATB I) and 180 mg day⁻¹ (ATB II). According to the *Fachinformation des Arzneimittel-Kompendium Schweiz*[®], the average oral dose (divided into three doses) of Clamoxyl[®]/- RC is 50 mg kg⁻¹ day⁻¹ for young children (2 to 12 year-old), with an absorption rate between 70 and 90% in the gastrointestinal tract. On the basis of two absorption rates of 70 and 85% we calculated that 90 to 180 mg day⁻¹ amoxicillin should reach the colon for a 12 kg-child receiving 600 mg amoxicillin per day. On the last evening of ATB I (day 25) and II (day 39), the reactor was briefly stopped and the medium was entirely pumped out of the reactor after beads had settled. A new medium without antibiotic (Stab II and III) was immediately added. Effluent samples (10 mL) were collected daily for metabolites (short-chain fatty acids [SCFA], and lactate) and fluorescent *in situ* hybridization (FISH) analyses. The pseudo-steady state for each period was reached when bacterial populations in the reactor

effluent did not change by more than 0.5 log units during 4 consecutive days (Cinquin et al., 2006).

During antibiotic treatments, effluent samples were collected before adding the first daily
170 antibiotic dose.

Bacterial enumeration with FISH and microscopy

FISH analyses coupled with microscopy were performed for total bacteria and *Salmonella* enumeration as described by Cinquin *et al.* (2006a) on fermentation samples (1.5 mL) from the last 3 days of each pseudo-steady-state period. Total bacteria were stained with 4', 6-
175 diamidino-2-phenylindole (DAPI, Sigma-Aldrich, Buchs, Switzerland) and *S. Typhimurium* was targeted with Sal 3, a Cy3-labelled oligonucleotide probe (Microsynth, Balgach, Switzerland) with hybridization conditions described by Nordentoft *et al.* (1997).

Bacterial enumeration with FISH and flow cytometry

FISH analyses coupled with flow cytometry were performed based on the method described
180 by Zoetendal *et al.* (2002) with some modifications. Briefly, 100 µL of fixed fermentation samples (1.5 mL) and fixed feces (1.5 mL) were centrifuged (9000 g, 3 min) and the pellet was washed once in Tris-EDTA buffer (100 mM Tris/HCl, 500 mM EDTA, pH 8) before incubation for 10 min at room temperature in 100 µL of Tris-EDTA buffer supplemented with lysozyme (170'800 U mL⁻¹) and proteinase K (6 µg mL⁻¹) to destroy protein-clusters formed
185 during both antibiotic treatment periods which interfered with flow cytometry detection. After removing the lysozyme solution by centrifugation (9000 g, 3 min) and washing the pellet once with 100 µL of fresh hybridization buffer (900 mM NaCl, 20 mM Tris/HCl, 0.1% SDS, 30% formamide, pH 8), cells were resuspended in 300 µL of hybridization buffer, homogenized and divided into 10 aliquots of 25 µL. With exception of the negative control,
190 aliquots were hybridized overnight at 35 °C with 50 ng µL⁻¹ of Cy5-labeled probes (Table 1). Since hybridization conditions described for Sal 3 (Nordentoft *et al.*, 1997) slightly differed

from conditions used in flow cytometry, the Sal 3 probe specificity was tested again with hybridization conditions used for flow cytometry (Table 1).

After hybridization and to remove non-specific binding of probes, 900 μL of warm washing

195 buffer (64 mM NaCl, 20 mM Tris/HCl, 5 mM EDTA, 0.1% SDS, pH 8) was added and samples were incubated at 37 °C for 20 min. A last centrifugation step was performed

(9000 g, 3 min) before resuspending the pellet in 300 μL of cold potassium citrate buffer

(10 mM Tris/HCl, 1 mM EDTA, 30 mM potassium citrate, pH 7.4). A 50 μL aliquote was

diluted with 400 μL of potassium citrate buffer to obtain a final bacterial concentration of

200 about 10^8 bacteria per mL. A volume of 0.5 μL of diluted (1/10'000) SYBR Green I

(Invitrogen AG, Basel, CH) was added at least 15 min before each measurement in order to

differentiate bacteria from non-bacterial material. To determine bacterial cell numbers, 50 μL

of Flow-CountTM Fluorospheres (Beckman Coulter International SA, Nyon, CH) at known

concentrations ($1012 \text{ beads } \mu\text{L}^{-1}$) were added just before data acquisition. Samples were

205 passed through a Cytomics FC 500 (Beckman Coulter International SA, Nyon, Switzerland)

equipped with an air-cooled argon ion laser emitting 20 mW at 488 nm and a Red Solid State

Diode laser emitting 25 mW at 633 nm. The 633 nm laser was used to detect red fluorescence

of bacteria hybridized with Cy5-labeled probes (PMT4 in a 655 nm long pass filter) and the

488 nm laser was used to measure the forward angle light scatter (FSC), the side angle light

210 scatter and the green fluorescence conferred by SYBR Green I (PMT1 in a 525 nm band pass

filter). The acquisition threshold was set in the forward scatter channel to the minimum. The

flow rate was set at $1,000\text{-}3,000 \text{ events s}^{-1}$ and 100,000 events were stored in list mode files.

Data were analyzed using the CXP software (Beckman Coulter International SA, Nyon,

Switzerland). A PMT1 histogram (green fluorescence) was used to evaluate the total number

215 of bacteria stained with SYBR Green I. In this histogram, a gate which included the total

number of bacterial cells in the sample was designed and used to make a PMT4 histogram

(red fluorescence). This PMT4 histogram was then used to determine the bacterial groups marked with Cy5-labelled probes. To quantify bacterial groups and total cells, a correction was made to eliminate background fluorescence, measured using the negative control NON-EUB338-Cy5 probe, as described by Rigottier-Gois *et al.* (2003). Analyses were done in duplicate.

Metabolite analyses

SCFA (acetate, propionate, butyrate and formate) and lactate concentrations were determined by HPLC as previously described (Cleusix *et al.*, 2008). Each analysis was done in duplicate.

Mean metabolite concentrations were expressed in mM.

Statistical analyses

A one-way ANOVA was performed using SPSS 13.0 for Windows (SPSS Inc., Chicago, IL, USA) to test the effects of the different treatments on bacterial and metabolite concentrations measured during the pseudo-steady-state periods (mean of 3 successive days) in effluent samples. Treatment means were compared using the Tukey's test with the probability level of $P < 0.05$. Data in the text are means \pm SD.

Results

Microbial populations analyzed by FISH-flow cytometry

The child fecal sample used for immobilization showed a total population of $10.2 \pm 0.1 \log_{10}$ cells g^{-1} , and was highly dominated by bifidobacteria ($9.8 \pm 0.1 \log_{10}$ cells g^{-1}). Mean concentrations of major bacterial populations measured in protease treated fermentation samples by FISH-Flow cytometry during the last three days of each experimental period are shown in Table 2. The predominant bacterial genus in the reactor effluents during the whole fermentation (except during STyphi I) was *Bifidobacterium* spp., initially followed by

240 *Clostridium coccooides-Eubacterium rectale* and *Bacteroides-Prevotella* groups. No
Salmonella or other *Enterobacteriaceae* were detected during STyphi 0. Following addition
 of *S. Typhimurium*-colonized beads ($11.1 \log_{10}$ cells (g bead)⁻¹) to the reactor (days 11 and
 13; STyphi I), a high concentration of *Salmonella* ($7.5 \pm 0.1 \log_{10}$ cells mL⁻¹) was measured in
 effluent samples at the end of the first stabilization period (STyphi I, Table 2). Furthermore,
 245 the intestinal microbial balance largely changed compared with STyphi 0, with a significant
 decrease of bifidobacteria (minus 1.2 log₁₀ unit) and a significant increase of
Enterobacteriaceae, *Atopobium* spp. and the *C. coccooides- E. rectale* group. During this
 period (STyphi I), the *C. coccooides- E. rectale* group became the predominant bacterial group.
 Addition of amoxicillin (90 mg day⁻¹, ATB I) induced a significant decrease in *Salmonella*
 250 concentration (more than 1.5 log₁₀ units) compared with STyphi I (Table 2) and a shift in the
 different bacterial concentrations, which returned to values not significantly different from
 STyphi 0 ($P > 0.05$). The only exception was *Enterobacteriaceae* which in contrast to
Salmonella, did not decrease during ATB I and remained significantly higher than during
 STyphi 0. During STyphi II with no antibiotic, most bacterial populations (including
 255 *Salmonella*) went back to values similar to STyphi I, except for *Bifidobacterium* spp. and *C.*
coccooides- E. rectale concentrations which were significantly higher and lower, respectively.
 The second amoxicillin treatment (180 mg day⁻¹, ATB II) induced similar changes as during
 ATB I, but the effects on bacterial populations measured during the last three days were
 significant only for *Salmonella* and the *C. coccooides- E. rectale* group. Finally, during the last
 260 stabilization period without antibiotic (STyphi III), bacterial populations went back to values
 similar to STyphi 0, except for *Salmonella* and total *Enterobacteriaceae* which remained
 significantly higher than for STyphi 0. No effect of treatments was observed on total bacteria
 or on *Bacteroides* concentrations during the whole fermentation.

Microbial populations analyzed by FISH-microscopy

265 Bacterial analyses with FISH coupled to microscopic counts for total bacteria and
S. Typhimurium used to monitor overtime the fermentation process corroborated data
obtained with flow cytometry, with a significant decrease in *Salmonella* concentration during
ATB I and II. No protease treatment was applied to these samples. However, a more
pronounced inhibition of *S. Typhimurium* was measured with ATB I (minus 1.3 log₁₀ units
270 compared with STyphi I) compared with ATB II (minus 0.95 log₁₀ units compared to
STyphi II) (Figure 2). Furthermore cell aggregation was enhanced during antibiotic
treatments. Total bacteria counts measured with FISH-microscopy were significantly lower
(minus 1.2 log₁₀ units) than by FISH-flow cytometry, decreased from STyphi 0 to STyphi I
and remained stable during the subsequent treatments.

275 ***Metabolites analyses***

The profiles of metabolite concentrations during the different fermentation periods are shown
in Figure 3. Acetate was the main metabolite detected during STyphi 0 and throughout the
fermentation (40-80 mM), whereas propionate and butyrate were present at low
concentrations (≤ 20 mM). Lactate was also detected, although in variable concentrations (1-
280 20 mM), during the whole fermentation. Addition of *S. Typhimurium* induced a significant
increase in butyrate (6.0 ± 0.1 and 13.5 ± 1.8 mM for STyphi 0 and STyphi I, respectively),
whereas the other metabolites were not modified between the two periods (Table 3). ATB I
dramatically decreased total metabolite concentration compared with STyphi I, especially due
to a strong decrease in acetate, while formate and lactate increased. During STyphi II,
285 metabolite concentrations went back to their previous levels of STyphi I, except for acetate
and butyrate which remained significantly lower than during STyphi I. ATB II induced
slightly different metabolite concentration changes compared with ATB I. A less pronounced
decrease in acetate concentration was observed, whereas lactate and formate were

significantly more increased. During the last stabilization period (STyphi III) metabolite concentrations returned to their previous levels of STyphi II, except for propionate and formate (Table 3).

Discussion

Recently we successfully developed a new *in vitro* model of intestinal fermentation with immobilized fecal microbiota (Cinquin *et al.*, 2004; Cinquin *et al.*, 2006a; Cinquin *et al.*, 2006b; Cleusix *et al.*, 2008). One major advantage of cell immobilization in intestinal fermentation models is the very high microbial and metabolic stability due to entrapment and growth of fecal microbiota in polysaccharide beads, which was tested over long fermentation periods (up to 7 weeks).

In this study, we used the same approach with fecal sample immobilization and continuous fermentation to develop an original model simulating *Salmonella* gut infection in children. We showed that cell immobilization can circumvent problems due to wash-out of exogenous enteropathogens observed in conventional *in vitro* intestinal fermentation models operated with planktonic cells (Blake *et al.*, 2003; Carman *et al.*, 2004; Payne *et al.*, 2003). According to our initial assumption, addition of *S. Typhimurium* immobilized in polysaccharide beads allowed to recover and maintain *S. Typhimurium* in the reactor effluents during the entire fermentation of 43 days. Seven days after addition of *S. Typhimurium* colonized beads (days 18-20; STyphi I), the strain was detected at high concentrations ($7.5 \log_{10}$ cells mL⁻¹) in effluent samples, in agreement with *in vivo* concentrations of non-typhoid *Salmonella* shedding of up to 10^6 - 10^7 organisms per gram of faeces measured in some children during early convalescence (Cruickshank & Humphrey, 1987). A daily analysis of *S. Typhimurium* and commensal bacteria was then performed in effluent samples. A protease treatment was used to destroy aggregates before bacterial enumeration with FISH-flow cytometry. However, no sample protease treatment was done before analysis with FISH-microscopy. Therefore the

aggregation phenomena can explain the lower total and *Salmonella* cell counts detected with
315 FISH-microscopy compared to FISH-flow cytometry. Furthermore the higher inhibition
effects of antibiotics on *Salmonella* detected with FISH-microscopy compared to FISH-flow
cytometry can also be due to enhanced cell aggregation observed during antibiotic treatments.
Total bacteria and bifidobacteria populations measured during STyphi 0 and in feces were
high and very close. Unfortunately, the other populations could not be determined in feces
320 due to a lack of sample. We showed in previous studies that the main populations of child and
adult fecal samples were well preserved during immobilization and long-term continuous
fermentation. However, differences of microbial balance between the fecal inoculum and
reactor effluents occurred due to changes of environmental conditions between the host
intestine and the fermentation model, such as medium composition, pH, and retention time
325 (Cinquin *et al.*, 2004; Cinquin *et al.*, 2006b, Cleusix *et al.*, 2008). Such differences are often
observed with *in vitro* intestinal fermentation systems (Macfarlane & Macfarlane, 2007).
Furthermore, conditions of the proximal colon applied in the fermentation model are very
different to that of the distal colon where composition of fecal material is more akin to feces.
Compared with STyphi 0, *S. Typhimurium* colonization (STyphi I) induced a strong
330 modification in the microbial balance. Bifidobacteria, present in high numbers during
STyphi 0 were significantly decreased, whereas *C. cocoides*- *E. rectale* group and *Atopobium*
spp. were strongly increased ($P < 0.05$). In contrast to bacterial populations, metabolites were
only slightly modified, with only a significant increase in butyrate concentrations during
STyphi I. This butyrate accumulation may be explained by an increase in the *C. cocoides*- *E.*
335 *rectale* group, which contains most of the butyrate producers (Barcenilla *et al.*, 2000).
As expected, addition of amoxicillin in the reactor significantly reduced *S. Typhimurium*
concentration in effluent samples and changed the microbial balance. *C. cocoides*- *E. rectale*
and *Atopobium* populations were strongly inhibited by amoxicillin, whereas *Bacteroides*,

bifidobacteria and total *Enterobacteriaceae* were not affected. Marked decreases in the *C. coccooides*- *E. rectale* group have already been described by Barc *et al.* (2004) in human fecal flora-associated mice receiving amoxicillin-clavulanic acid (150 mg (kg of body weight)⁻¹) for 7 days. The lack of activity of amoxicillin on *Bacteroides-Prevotella* and *Enterobacteriaceae* was also expected because it is known that certain *Bacteroides* such as *Bacteroides fragilis* and some *Enterobacteriaceae* (*Klebsiella* spp., *E. coli*) are not inhibited during amoxicillin treatments in humans (Sullivan *et al.*, 2001; Floor *et al.*, 1994). Indeed, resistance to β -lactams via β -lactamase production has been largely described in *Bacteroides* spp. and *Enterobacteriaceae* (Kader *et al.*, 2004; Papaparaskevas *et al.*, 2005). Unexpectedly, bifidobacteria were not inhibited by amoxicillin treatments and their growth was even stimulated during ATB I. Although it is generally admitted that bifidobacteria are highly sensitive to β -lactams (Moubareck *et al.*, 2005) some strains are resistant to amoxicillin (Lim *et al.*, 1993; Vlková *et al.*, 2006). However, bifidobacteria overgrowth during ATB I may not be directly associated with the effects of antibiotics. It could also be due to the strong decrease in the *C. coccooides*- *E. rectale* group, which allowed the growth of bifidobacteria that came back to their previous concentrations (STyphi 0). It seems that bifidobacteria were particularly competitive in the system and that they prevented the overgrowth of *Bacteroides* spp. during amoxicillin treatment, contrary to some *in vivo* observations (Christensson *et al.*, 1991). *Enterobacteriaceae* were not significantly decreased during ATB I, and ATB II despite a significant decrease of *Salmonella*, and became the second most dominant group during STyphi II, taking the place of the *C. coccooides*- *E. rectale* group. Such increases in *Enterobacteriaceae* have also been frequently described in the literature after amoxicillin treatments in humans (Christensson *et al.*, 1991; Sullivan *et al.*, 2001).

In parallel to modifications of microbial balance during ATB I and II, metabolite ratios were also strongly modified. ATB I decreased acetate and increased lactate and formate concentrations, whereas butyrate and propionate were not changed compared with STyphi I. Doubling the amoxicillin concentration (ATB II) also led to a similar decrease in acetate concentration compared with STyphi II, but also to twice as much lactate and formate and to a decrease in propionate and butyrate concentrations. Lactate and formate are intermediate metabolites produced by many different bacteria (Cummings & Macfarlane, 1991), further metabolised by other bacteria into CO₂ and major SCFA (acetate, propionate or butyrate) (Duncan *et al.*, 2004; Seeliger *et al.*, 2002). These metabolites generally do not accumulate in the human colon (Bernalier *et al.*, 1999). Their accumulation together with the changes in SCFA ratios during ABT I and II suggest that lactate- and formate-utilizing bacteria were inactivated during amoxicillin treatments. The marked decrease in the *C. cocoides-E. rectale* group, which harbours many lactate-utilizing bacteria (Duncan *et al.*, 2004) is likely responsible for lactate accumulation during these periods.

Immobilization of *S. Typhimurium* and its addition to the recently developed *in vitro* model of intestinal fermentation with immobilized fecal microbiota led to stable high levels of *S. Typhimurium* in the effluent of the continuous gut reactor simulating *Salmonella* shedding observed in certain children. Moreover, immobilization prevented washout of *S. Typhimurium* during antibiotic treatments, which allowed comparison of two treatments during the same fermentation with the same fecal inoculum. The effects of amoxicillin measured with this new *in vitro* colonic fermentation model for *Salmonella* infections are in agreement with *in vivo* observations showing a high disturbance of the intestinal microbiota balance with decreases in the *C. coccooides* - *E. rectale* group and increases in *Enterobacteriaceae* ratios. Furthermore, metabolic ratios tested in the *in vitro* model correlated with microbial change, providing further validation to this model. Therefore, this

new model is a promising tool for simulating intestinal infections in humans with the aim of developing and testing the effects of different antimicrobials on intestinal enteropathogens as well as commensal bacteria.

Acknowledgements

This research was supported by a grant from the Swiss National Science Foundation (SNF) (project number: 3100170-114028).

References

- 395 Barc MC, Bourlioux F, Rigottier-Gois L, Charrin-Sarnel C, Janoir C, Boureau H, Doré J, Collignon A (2004) Effect of amoxicillin-clavulanic acid on human fecal flora in a gnotobiotic mouse model assessed with fluorescence hybridization using group-specific 16S rRNA probes in combination with flow cytometry. *Antimicrob Agents Chemother* **48**: 1365-1368.
- 400 Barcenilla A, Pryde SE, Martin JC, Duncan SH, Stewart CS, Henderson C, Flint HJ (2000) Phylogenetic relationships of butyrate-producing bacteria from the human gut. *Appl Environ Microbiol* **66**: 1654-1661.
- Bernalier A, Dore J, Durand M (1999) Biochemistry of fermentation. In: Gibson GR and Roberfroid MB (Ed.). *Colonic microbiota, nutrition and health*. Kluwer academic publishers, London. pp 37-53.
- 405 Blake DP, Hillman K, Fenlon DR (2003) The use of a model ileum to investigate the effects of novel and existing antimicrobials on indigenous porcine gastrointestinal microflora: using vancomycin as an example. *Anim Feed Sci Tech* **103**: 123-139.
- Carman RJ, Simon MA, Fernandez H, Miller MA, Bartholomew MJ (2004) Ciprofloxacin at low levels disrupts colonization resistance of human fecal microflora growing in chemostats. *Regul Toxicol Pharmacol* **40**: 319-326.
- 410 Carman RJ & Woodburn MA (2001) Effects of low levels of ciprofloxacin on a chemostat model of the human colonic microflora. *Regul Toxicol Pharm* **33**: 276-284.
- Cebrian L, Rodriguez JC, Escribano I, Royo G (2006) *In vitro* generation of *Salmonella* spp. mutants following repeated exposure to beta-lactam antibiotics. *J Infect Chemother* **12**: 80-82.
- 415

Christensson B, Nilsson-Ehle I, Ljungberg B, Nömm I, Oscarsson G, Nordström L, Goscinsky G, Löwdin E, Linglöv T, Nordström B *et al.* (1991) A randomized multicenter trial to compare the influence of cefaclor and amoxycillin on the colonization resistance of the digestive tract in patients with lower respiratory tract infection. *Infection* **19**: 208-15.

420 Cinquin C, Le Blay G, Fliss I, Lacroix C (2004) Immobilization of infant fecal microbiota and utilization in an *in vitro* colonic fermentation model. *Microb Ecol* **48**: 128-138.

Cinquin C, Le Blay G, Fliss I, Lacroix C (2006a) Comparative effects of exopolysaccharides from lactic acid bacteria and fructo-oligosaccharides on infant gut microbiota tested in an *in vitro* colonic model with immobilized cells. *FEMS Microbiol Ecol* **57**: 226-238.

425 Cinquin C, Le Blay G, Fliss I, Lacroix C (2006b) New three-stage *in vitro* model for infant colonic fermentation with immobilized fecal microbiota. *FEMS Microbiol Ecol* **57**: 324-336.

Cleusix V, Lacroix C, Vollenweider S, Le Blay G (2008) Glycerol induces reuterin production and decreases *Escherichia coli* population in an *in vitro* model of colonic fermentation with immobilized human feces. *FEMS Microbiol Ecol* **63**: 56-64.

430 Coburn B, Grassl GA, Finlay BB (2007) *Salmonella*, the host and disease: a brief review. *Immunol Cell Biol* **85**: 112-118.

Cruickshank JG & Humphrey TJ (1987) The carrier food-handler and non-typhoid salmonellosis. *Epidemiol Infect* **98**: 223-230.

Cummings JH & Macfarlane GT (1991) The control and consequences of bacterial
435 fermentation in the human colon. *J Appl Bacteriol* **70**: 443-459.

<http://www.kompendium.ch/Monographie.aspx?Id=8553f7cf-bda5-4841-baed-86f3a7e32b27&lang=de&MonType=fi>

440 Duncan SH, Louis P, Flint HJ (2004) Lactate-utilizing bacteria, isolated from human feces,
that produce butyrate as a major fermentation product. *Appl Environ Microbiol* **70**: 5810-
5817.

Fallingborg J, Christensen LA, Ingeman-Nielsen M, Jacobsen BA, Abildgaard K, Rasmussen
HH, Rasmussen SN (1990) Measurement of gastrointestinal pH and regional transit times in
445 normal children. *J Pediatr Gastroenterol Nutr* **11**: 211-214.

Floor M, van Akkeren F, Rozenberg-Arska M, Visser M, Kolsters A, Beumer H, Verhoef J
(1994) Effect of loracarbef and amoxicillin on the oropharyngeal and intestinal microflora of
patients with bronchitis. *Scand J Infect Dis* **26**: 191-197.

Franks AH, Harmsen HJ, Raangs GC, Jansen GJ, Schut F, Welling GW (1998) Variations of
450 bacterial populations in human feces measured by fluorescent *in situ* hybridization with
group-specific 16S rRNA-targeted oligonucleotide probes. *Appl Environ Microbiol* **64**: 3336-
3345.

Frye JG & Fedorka-Cray PJ (2007) Prevalence, distribution and characterisation of ceftiofur
resistance in *Salmonella enterica* isolated from animals in the USA from 1999 to 2003. *Int J*
455 *Antimicrob Agents* **30**: 134-142.

Hapfelmeier S, Ehrbar K, Stecher B, Barthel M, Kremer M, Hardt WD (2004) Role of the
Salmonella pathogenicity island 1 effector proteins SipA, SopB, SopE, and SopE2 in

Salmonella enterica subspecies 1 serovar Typhimurium colitis in streptomycin-pretreated mice. *Infect Immun* **72**: 795-809.

460 Hapfelmeier S & Hardt WD (2005) A mouse model for *S. typhimurium*-induced enterocolitis. *Trends Microbiol* **13**: 497-503.

Harmsen HJM, Wildeboer-Veloo ACM, Grijpstra J, Knol J, Degener JE, Welling GW (2000) Development of 16S rRNA-based probes for the *Coriobacterium* Group and the *Atopobium* cluster and their application for enumeration of *Coriobacteriaceae* in human feces from
465 volunteers of different age groups. *Appl Environ Microbiol* **66**: 4523-4527.

Kader AA, Kumar A, Dass SM (2004) Antimicrobial resistance patterns of Gram-negative bacteria isolated from urine cultures at a general hospital. *Saudi J Kidney Dis Transpl* **15**: 135-139.

Kiessling CR, Jackson M, Watts KA, Loftis MH, Kiessling WM, Buen MB, Laster EW,
470 Sofos JN (2007) Antimicrobial susceptibility of *Salmonella* isolated from various products, from 1999 to 2003. *J Food Prot* **70**: 1334-1338.

Langendijk PS, Schut F, Jansen GJ, Raangs GC, Kamphuis GR, Wilkinson MH, Welling GW (1995) Quantitative fluorescence *in situ* hybridization of *Bifidobacterium* spp. with genus-specific 16S rRNA-targeted probes and its application in fecal samples. *Appl Environ*
475 *Microbiol* **61**: 3069-3075.

Lim KS, Huh CS, Baek YJ (1993) Antimicrobial susceptibility of bifidobacteria. *J Dairy Sci* **76**: 2168-2174.

- Macfarlane GT & Macfarlane S (2007) Models for intestinal fermentation: association
between food components, delivery systems, bioavailability and functional interactions in the
480 gut. *Current Opin Biotech* **18**: 156-162.
- Macfarlane GT, Macfarlane S, Gibson GR (1998) Validation of a three-stage compound
continuous culture system for investigating the effect of retention time on the ecology and
metabolism of bacteria in the human colon. *Microb Ecol* **35**: 180-187.
- Macfarlane S, McBain AJ, Macfarlane GT (1997). Consequences of biofilm and sessile
485 growth in the large intestine. *Adv Dent Res* **11**: 59-68.
- Manz W, Amann R, Ludwig W, Vancanneyt M, Schleifer KH (1996) Application of a suite of
16S rRNA-specific oligonucleotide probes designed to investigate bacteria of the phylum
cytophaga-flavobacter-bacteroides in the natural environment. *Microbiology* **142**: 1097-1106.
- Michel C, Kravtchenko TP, David A, Gueneau S, Kozlowski F, Cherbut C (1998) *In vitro*
490 prebiotic effects of Acacia gums onto the human intestinal microbiota depends on both origin
and environmental pH. *Anaerobe* **4**: 257-266.
- Moubareck C, Gavini F, Vaugien L., Butel MJ, Doucet-Populaire F (2005). Antimicrobial
susceptibility of bifidobacteria. *J Antimicrob Chemother* **55**: 38-44.
- Moulin F, Sauve-Martin H, Marc E, Lorrot MM, Soulier M, Ravilly S, Raymond J, Gendrel D
495 (2003) Ciprofloxacin after clinical failure of beta-lactam antibiotics in children with
salmonellosis. *Arch Pediatr* **10**: 608-14.
- Nordentoft S, Christensen H, Wegener HC (1997) Evaluation of a fluorescence-labelled
oligonucleotide probe targeting 23S rRNA for *in situ* detection of *Salmonella* serovars in

- paraffin-embedded tissue sections and their rapid identification in bacterial smears. *J Clin Microbiol* **35**: 2642-2648.
- Papaparaskevas J, Pantazatou A, Katsandri A, Legakis NJ, the Hellenic Study Group for Gram-Negative Anaerobic Bacteria, Avlami A (2005) Multicentre survey of the *in vitro* activity of seven antimicrobial agents, including ertapenem, against recently isolated Gram-negative anaerobic bacteria in Greece. *Clin Microbiol Infect* **11**: 820-824.
- 505 Park ES, Park CI, Cho SR, Na SI, Cho YS (2004) Colonic transit time and constipation in children with spastic cerebral palsy. *Arch Phys Med Rehabil* **85**: 453-456.
- Payne S, Gibson G, Wynne A, Hudspith B, Brostoff J, Tuohy K (2003) *In vitro* studies on colonization resistance of the human gut microbiota to *Candida albicans* and the effects of tetracycline and *Lactobacillus plantarum* LPK. *Curr Issues Intest Microbiol* **4**: 1-8.
- 510 Rigottier-Gois L, Le Bourhis A-G, Gramet G, Rochet V, Doré J (2003) Fluorescent hybridisation combined with flow cytometry and hybridisation of total RNA to analyse the composition of microbial communities in human faeces using 16S rRNA probes. *FEMS Microb Ecol* **43**: 237-245.
- Rosanova MT, Paganini H, Bologna R, Lopardo H, Ensinn G (2002) Risk factors for mortality caused by nontyphoidal *Salmonella* sp. in children. *Int J Infect Dis* **6**: 187-190.
- 515 Santos RL, Tsois RM, Bäumlér AJ, Adams LG (2003) Pathogenesis of *Salmonella*-induced enteritis. *Braz J Med Biol Res* **36**: 3-12.
- Schaad UB (2005) Fluoroquinolone antibiotics in infants and children. *Infect Dis Clin North Am* **19**: 617-28.

- 520 Seeliger S, Janssen PH, Schink B (2002) Energetics and kinetics of lactate fermentation to acetate and propionate via methylmalonyl-CoA or acrylyl-CoA. *FEMS Microbiol Lett* **211**: 65-70.
- Sghir A, Gramet G, Suau A, Rochet V, Pochart P, Dore J (2000) Quantification of bacterial groups within human fecal flora by oligonucleotide probe hybridization. *Appl Environ*
- 525 *Microbiol* **66**: 2263-2266.
- Stark CA, Edlund C, Sjöstedt S, Kristensen G, Nord CE (1993) Antimicrobial resistance in human oral and intestinal anaerobic microfloras. *Antimicrob Agents Chemother* **37**: 1665-1669.
- Sullivan A, Edlund C, Nord CE (2001). Effect of antimicrobial agents on the ecological
- 530 balance of human microflora. *Lancet Infect Dis* **1**: 101-114.
- The European Food Safety Authority & European Center for Disease Prevention and Control (2006) The community summary report on trends and sources of zoonoses, zoonotic agents, antimicrobial resistance and foodborne outbreaks in the European Union in 2005. *The EFSA journal* **94**: 1-236.
- 535 Vlková E, Rada V, Popelářřová P, Trojanová I, Killer J (2006) Antimicrobial susceptibility of bifidobacteria isolated from gastrointestinal tract of calves. *Livestock Science* **105**: 253-259.
- Wallner G, Amann R, Beisker W (1993) Optimizing fluorescent *in situ* hybridization with ribosomal-RNA-targeted oligonucleotide probes for flow cytometric identification of microorganisms. *Cytometry* **14**: 136-143.

540 White DG, Zhao S, Sudler R, Ayers S, Friedman S, Chen S, McDermott PF, McDermott S,
Wagner DD, Meng J (2001). The isolation of antibiotic-resistant *Salmonella* from retail
ground meats. *N Engl J Med* **345**: 1147-1154.

World Health Organisation, © Copyright by World Health Organization (2008) Drug-resistant
Salmonella (Fact sheet N°139)

545 <http://www.who.int/mediacentre/factsheets/fs139/en/index.html>

Zoetendal EG, Ben-Amor K, Harmsen HJ, Schut F, Akkermans AD, de Vos WM (2002)
Quantification of uncultured *Ruminococcus obeum*-like bacteria in human fecal samples by
fluorescent in situ hybridization and flow cytometry using 16S rRNA-targeted probes. *Appl*
Environ Microbiol **68**: 4225-4232.

550

Tables

Table 1 Oligonucleotide probes and hybridization conditions used to target predominant bacterial groups

Probes	Dyes*	Target organisms	Formamide (%)	Temperature (°C)	Probe references
Sal 3	Cy-3	<i>Salmonella enterica</i>	0	45	Nordentoft <i>et al.</i> , 1997
NonEub338	Cy-5	NA	30	35	Wallner <i>et al.</i> , 1993
Bif 164	Cy-5	<i>Bifidobacterium</i>	30	35	Langendijk <i>et al.</i> , 1995
Bac 303	Cy-5	<i>Bacteroides-Prevotella</i>	30	35	Manz <i>et al.</i> , 1996
Erec 482	Cy-5	<i>C. coccoides-E. rectale</i>	30	35	Franks <i>et al.</i> , 1998
Ato 291	Cy-5	<i>Atopobium</i>	30	35	Harmsen <i>et al.</i> , 2000
Enter 432	Cy-5	Enterobacteria	30	35	Sghir <i>et al.</i> , 2000
Sal 3	Cy-5	<i>Salmonella</i>	30	35	Nordentoft <i>et al.</i> , 1997

*The oligonucleotide probe labelled at the 5' end with Cy-3 was detected with microscopy whereas Cy-5 labelled probes were used in flow cytometry. DAPI and SYBR Green I-stains were used for total bacteria enumeration in microscopy and flow cytometry, respectively.

Table 2 Bacterial populations in fermentation samples during pseudo-steady states of each treatment period measured by FISH-flow cytometry

Bacterial populations	Bacterial counts (\log_{10} cells mL^{-1} medium)*					
	STyphi 0	STyphi I	ATB I	STyphi II	ATB II	STyphi III
<i>Bifidobacterium</i>	$10.2 \pm 0.1\text{c}$	$9.0 \pm 0.2\text{a}$	$10.5 \pm 0.3\text{c}$	$9.9 \pm 0.3\text{bc}$	$9.7 \pm 0.5\text{ab}$	$10.3 \pm 0.1\text{c}$
<i>C. coccoides-E. rectale</i>	$9.1 \pm 0.1\text{bc}$	$10.3 \pm 0.3\text{d}$	$8.5 \pm 0.1\text{ab}$	$9.5 \pm 0.3\text{c}$	$8.4 \pm 0.4\text{a}$	$8.5 \pm 0.4\text{ab}$
<i>Bacteroides</i>	$8.8 \pm 0.2\text{a}$	$9.0 \pm 0.2\text{a}$	$9.0 \pm 0.5\text{a}$	$9.0 \pm 0.2\text{a}$	$8.6 \pm 0.1\text{a}$	$9.0 \pm 0.02\text{a}$
<i>Atopobium</i>	$7.3 \pm 0.04\text{a}$	$9.2 \pm 0.1\text{c}$	$7.5 \pm 0.4\text{a}$	$8.7 \pm 0.3\text{bc}$	$8.0 \pm 0.5\text{ab}$	$7.2 \pm 0.7\text{a}$
<i>Enterobacteriaceae</i>	NDc	$8.9 \pm 0.2\text{b}$	$8.2 \pm 0.3\text{ab}$	$8.3 \pm 0.2\text{ab}$	$7.4 \pm 0.3\text{a}$	$9.4 \pm 0.3\text{b}$
<i>Salmonella</i>	ND	$7.5 \pm 0.1\text{a}$	NDb	$8.1 \pm 0.2\text{a}$	NDb	$7.6 \pm 0.2\text{a}$
Total bacteria	$10.3 \pm 0.4\text{a}$	$10.4 \pm 0.2\text{a}$	$10.5 \pm 0.1\text{a}$	$10.3 \pm 0.1\text{a}$	$10.0 \pm 0.3\text{b}$	$10.4 \pm 0.2\text{a}$

*Data are means \pm S.D. for the last 3 days for each fermentation period, $n = 2$. Values with different letters in a row are significantly different with the Tukey's test, $P < 0.05$.

[†]ND, not detected, below detection limit of the method ($\log_{10} 6$ cells mL^{-1}).

Table 3 Metabolite concentrations in effluent samples during pseudo-steady states of each treatment period measured by HPLC

Metabolites	Metabolite concentration (mM)*					
	STyphi 0	STyphi I	ATB I	STyphi II	ATB II	STyphi III
Acetate	69.6 ± 2.1bc	72.5 ± 3.6c	45.9 ± 5.1a	63.6 ± 2.7b	50.1 ± 3.7a	64.8 ± 1.7bc
Propionate	5.2 ± 0.5b	6.1 ± 2.0bc	6.2 ± 0.6bc	7.5 ± 1.0c	2.2 ± 0.4a	3.8 ± 0.2ab
Butyrate	6.0 ± 0.1ab	13.5 ± 1.8c	13.3 ± 2.6c	8.2 ± 1.5b	3.9 ± 1.5a	7.9 ± 1.5ab
Formate	NDa [†]	NDa	3.8 ± 0.8b	NDa	7.2 ± 1.8bc	10.5 ± 1.4c
Lactate	1.3 ± 0.2a	1.2 ± 0.4a	7.9 ± 1.0b	2.0 ± 0.5a	17.1 ± 3.0c	4.1 ± 1.6d
Total metabolites	82.1 ± 1.6abc	93.3 ± 4.0c	77.1 ± 4.2a	81.3 ± 2.4abc	80.5 ± 5.1ab	91.2 ± 3.5bc

*Data are means ± S.D. for the last 3 days for each fermentation period, $n = 2$. Values with different letters in a row are significantly different with the Tukey's test, $P < 0.05$.

[†]ND, not detected, below detection limit of the method.

Figure legends

Fig. 1. Time schedule of continuous intestinal fermentation experiments during different treatment periods: BC, beads colonization; STyphi 0, stabilization of the system, days 3-10; SB, addition of *S. Typhimurium* M557 beads, days 11 and 13; STyphi I, stabilization of *S. Typhimurium* M557 concentrations in effluent samples, days 14-20; ATB I, daily addition of 90 mg amoxicillin, days 21-25; STyphi II, stabilization period, days 26-34; ATB II, daily addition of 180 mg amoxicillin, days 35-39, STyphi III; stabilization period, days 40-43.

Fig. 2. Total bacteria and *Salmonella* enumerated by FISH coupled with microscopy. DAPI and Cy3 labeled Sal 3-probe were used to detect total bacteria and *Salmonella*, respectively, during the different fermentation periods. STyphi 0, stabilization period I without *Salmonella*. STyphi I, stabilization period II after *Salmonella* addition; ATB I, first antibiotherapy, 90 mg day⁻¹; STyphi II, stabilization period; ATB II, second antibiotherapy, 180 mg day⁻¹; STyphi III, stabilization period. Significance is based on last three days of each period.

Fig. 3. Short chain fatty acids and lactate concentrations in effluent samples during the 43-day continuous fermentation. Total metabolites (◆), acetate(●), butyrate (Δ), propionate (▲), formate (○), lactate (◇). Data are means of triplicate analyses.

Figure 1




Figure 2




Figure 3

